

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
12 August 2004 (12.08.2004)

PCT

(10) International Publication Number
WO 2004/068116 A2

(51) International Patent Classification⁷:

G01N

(21) International Application Number:

PCT/US2004/000425

(22) International Filing Date: 9 January 2004 (09.01.2004)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

60/440,840	17 January 2003 (17.01.2003)	US
60/440,838	17 January 2003 (17.01.2003)	US
10/379,965	4 March 2003 (04.03.2003)	US

(71) Applicant (for all designated States except US): ACLARA BIOSCIENCES, INC. [US/US]; 1288 Pear Avenue, Mountain View, CA 94043 (US).

(72) Inventor; and

(75) Inventor/Applicant (for US only): SINGH, Sharat [US/US]; 27359 Julietta Lane, Los Altos Hills, CA 94022 (US).

(74) Agent: MACEVICZ, Stephen; Aclara Biosciences, Inc., 1288 Pear Avenue, Mountain View, CA 94043 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to the identity of the inventor (Rule 4.17(i)) for all designations
- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for all designations
- of inventorship (Rule 4.17(iv)) for US only

Published:

- without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: RECEPTOR DEORPHANIZATION USING TAGGED MOLECULAR LIBRARIES

(57) Abstract: The present invention provides a method of screening test compounds and test conditions for the ability to modulate (activate or inhibit, enhance or depress) a GPCR pathway, and provides methods of assessing GPCR pathway function, such as the function of an orphan GPCR, in a cell in general. In another aspect of the present method, lipophilic photosensitizers are attached to the cellular membranes. A candidate ligand or a library of candidate ligands can be attached to a molecular, after which the ligand is allowed to bind to the receptor. After excitation of the photosensitizer with a light source, the cleavable linker is cleaved releasing the molecular tag. The released molecular tag can be detected in the extracellular fluid, as detailed above, which provides information on the structure of the ligand for the GPCR.